## **SURGERY POST OPERATIVE INSTRUCTIONS**



Some degree of pain and swelling are considered normal. Skin discolorations may also be experienced. Maximum swelling will occur between 2-3 days after the procedure. It is important that these instructions are followed to allow uncomplicated and rapid healing.

## Day 1:

- 1. A small amount of bleeding is to be expected following the operation. Place a gauze pad between your cheek and the tooth and maintain the pressure. Replace every 30 minutes as needed. Expect some oozing for the first 24 hours
- 2. Apply an ice pack to face over the operated area 20 minutes on and 20 minutes off until bedtime in order to keep the swelling to a minimum. A bag of frozen peas placed in Ziploc bags works well
- 3. Do not raise your lip with your fingers to inspect the treated area. The operated area should be uninterrupted to facilitate healing
- 4. Rest for the entire day
- 5. Only COLD liquids are allowed. Avoid hot liquids for the first 48 hours
- 6. Refrain from brushing and rinsing
- 7. Refrain from smoking for 12 hours
- 8. Elevate your head on a few pillows while sleeping to keep swelling to a minimum
- 9. Take medications as prescribed. The prescribed pain medicine may make you groggy and may slow down your reflexes. Do not drive an automobile or work around machinery. Avoid alcoholic beverages

## Day 2-7:

- 1. Discontinue the applications of ice to the face
- 2. Begin to rinse the treated area with warm saltwater 5-8 times daily. Use about ½ teaspoon of salt to 8 oz. of warm water. Make sure you soak the area each time for about 5 minutes
- 3. Brushing your teeth is allowed EXCEPT for the area that was treated surgically. You may also rinse your mouth after brushing and after meals to keep the surgical area clean
- 4. Get plenty of rest and insure normal intake of food, especially liquids such as fruit juices, soups and milk.
- 5. Resume exercise after the third day
- 6. Take medications as prescribed. Pain or discomfort following surgery should subside more and more every day. If the pain does not begin to subside in 2–3 days, or increases after 3 days, please call our office.
- 7. If sutures are placed, we will see you in our office in 5-7 days to follow and remove the sutures

Should any difficulties or questions arise, do not hesitate to text or call our office (281) 961-0961